



Walgreens

-PE Product Labels –

-continued



NDC 0363-0763-40

Walgreens

Compare to the active ingredients in Vicks[®] NyQuil[®] Severe Cold & Flu^{††}

NIGHTTIME

Severe Cold & Flu

ACETAMINOPHEN /
ACHES / FEVER / SORE THROAT
DEXTROMETHORPHAN HBr / COUGH
DOXYLAMINE SUCCINATE /
SNEEZING / RUNNY NOSE
PHENYLEPHRINE HCl / NASAL & SINUS
CONGESTION / SINUS PRESSURE

Maximum Strength

- Pain reliever, fever reducer, cough suppressant, antihistamine & nasal decongestant
- Alcohol free

Mixed Berry flavor

12 FL OZ (355 mL)

: 76340 94 F9

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.

ITEM 330480 W10107-0622-F
W3ORG0222-F

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Gluten Free

DO NOT USE IF PRINTED
NECKBAND IS BROKEN OR MISSING

3 11917 16341 3

Drug Facts

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine
Phenylephrine HCl 10 mg.....	Nasal decongestant

Uses: temporarily relieves common cold/flu symptoms: ■ sinus congestion and pressure ■ nasal congestion ■ minor aches and pains ■ headache ■ runny nose and sneezing ■ sore throat ■ cough to help you sleep ■ fever ■ cough due to minor throat and bronchial irritation ■ reduces swelling of nasal passages ■ promotes nasal and/or sinus drainage ■ temporarily restores freer breathing through the nose

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

PEEL BACK AT CORNER FOR MORE INFORMATION

: 76340 94 B1

Drug Facts
(continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have ■ liver disease ■ heart disease

WESIVE AREA
• NO VARNISH • NO TYPE

Drug Facts
(continued)

When using this product ■ do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ take only as directed – see Overdose warning ■ only use the dose cup

ADHESIVE AREA
ING • NO VARNISH • NO TYPE

AD
NO COATING

- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

◀

3177

provided. **do not exceed 4 doses per 24 hrs**

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information ■ each 30 mL contains: sodium 41 mg ■ store at 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions?
1-800-719-9260

*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
†This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks® and NyQuil®.

NO COAT



CONVENIENT RECLOSEING TAB

DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED

Drug Facts

Active ingredients (in each packet)

Acetaminophen 650 mg.....	Pain reliever/fever reducer
Diphenhydramine HCl 25 mg.....	Antihistamine/cough suppressant
Phenylephrine HCl 10 mg.....	Nasal decongestant

Purposes

Uses

- temporarily relieves these symptoms due to a cold:
 - minor aches and pains
 - minor sore throat pain
 - headache
 - nasal and sinus congestion
 - runny nose
 - sneezing
 - itchy nose or throat
 - itchy, watery eyes due to hay fever
 - cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not exceed recommended dosage
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not use more than directed (see overdose warning)

- take every 4 hours, while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and over	one packet
children under 12 years of age	do not use

- dissolve contents of one packet into 8 oz. hot water: sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water: stir briskly before and after heating. Do not overheat.

Other information

- each packet contains: potassium 10 mg and sodium 25 mg
- phenylketonurics: contains phenylalanine 13 mg per packet
- store at 20-25°C (68-77°F). Protect product from heat and moisture.

Inactive ingredients

acesulfame potassium, anhydrous citric acid, aspartame, colloidal silicon dioxide, D&C yellow #10, FD&C blue #1, FD&C red #40, flavors, maltodextrin, pregelatinized starch, sodium citrate, sucrose, tribasic calcium phosphate

Questions or comments?


1-800-719-9260

Gluten Free

OPEN OTHER END

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.
MADE IN MEXICO

ITEM 538582 W10106-0522-L



Walgreens



DAY & NIGHT PACK

NDC 0302-1507-02

Walgreens

Compare to the active ingredients in
Vicks® DayQuil® Severe Cold & Flu
& Vicks® NyQuil® Severe Cold & Flu†

DAYTIME
NON-DROWSYSevere
Cold
& Flu

ACETAMINOPHEN / ACHES /
FEVER / SORE THROAT
DEXTROMETHORPHAN HBr / COUGH
GUAIFENESIN / CHEST CONGESTION
PHENYLEPHRINE HCl /
NASAL CONGESTION

Maximum Strength

- Pain reliever, fever reducer,
cough suppressant, expectorant
& nasal decongestant
- Antihistamine free
- Alcohol free

NIGHTTIME

Severe
Cold
& Flu

ACETAMINOPHEN / ACHES /
FEVER / SORE THROAT
DEXTROMETHORPHAN HBr / COUGH
DOXYLAMINE SUCCINATE /
SNEEZING / RUNNY NOSE
PHENYLEPHRINE HCl /
NASAL & SINUS CONGESTION /
SINUS PRESSURE

Maximum Strength

- Pain reliever, fever reducer, cough
suppressant, antihistamine
& nasal decongestant
- Alcohol free

Mixed Berry
flavor

2 - 12 FL OZ (355 mL) BOTTLES / TOTAL 24 FL OZ (1.5 pt) (710 mL)

DO NOT USE IF PRINTED NECKBAND IS
BROKEN OR MISSING

Gluten Free



how2recycle.info

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

†Our pharmacists recommend the
Walgreens brand. We invite you to
compare to national brands.
‡These products are not manufactured or
distributed by The Procter & Gamble
Company, owner of the registered
trademarks Vicks® NyQuil® and DayQuil®.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015

Walgreens
100% SATISFACTION
GUARANTEED

ITEM 957097 WI 0108-0722-L



MightyVive Severe Cold & Flu**Drug Facts****Active Ingredients (in each 30 mL)**

Acetaminophen 160 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine
Phenylephrine HCl 10 mg	Nasal decongestant

Purpose**Uses** temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- headache
- fever
- sore throat
- runny nose and sneezing
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
 - with other drugs containing acetaminophen
 - 2 or more alcoholic drinks every day while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough/asthma occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking antibiotics or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Take only as directed – see Overdose warning

- only use the dose cup provided
 - do not exceed 4 doses per 24 hrs
- | | |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 4 to under 12 yrs | ask a doctor |
| children under 4 yrs | do not use |

Other information

- each 30 mL contains: sodium 41 mg
- store at 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments? 1-800-716-8286

MightyVive Severe Cold & Flu**Drug Facts****Active Ingredients (in each 15 mL)**

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Purpose**Uses** temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passages of both mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 4,000 mg of acetaminophen in 24 hours
 - child takes more than 5 doses in 24 hours
 - takes with other drugs containing acetaminophen
 - adult has 2 or more alcoholic drinks every day while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**When using this product do not use more than directed****Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Take only as directed – see Overdose warning

- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 8 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients buffered hydroxyanisole, edetate disodium, FD&C yellow #6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments? 1-800-716-8286

W3ORG0422-F

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not use more than directed

Stop use and ask a doctor if ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ you get nervous, dizzy or sleepless ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ take only as directed – see Overdose warning
■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information ■ each 30 mL contains: sodium 41 mg
■ store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

Questions or comments? 1-800-719-9260

Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey
† This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® DayQuil® Severe+ VapoCOOL™.

ITEM 542719 W10107-0719-F

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

3 11917 20841 1

Walgreens

Compare to Vicks® DayQuil® Severe+ VapoCOOL™ active ingredients!

DAYTIME • NON-DROWSY

Severe Cold & Flu

ACETAMINOPHEN / MINOR ACHES & PAINS / FEVER
PHENYLEPHRINE HCl / NASAL CONGESTION / SINUS PRESSURE
DEXTROMETHORPHAN HBr / COUGH
GUAIFENESIN / CHEST CONGESTION

NDC 0363-1940-40

NEW

MAXIMUM STRENGTH

- Pain reliever, fever reducer, cough suppressant, expectorant, nasal decongestant
- ALCOHOL 10%

12 FL OZ (355 mL)

VAPOR ICE™

Drug Facts **DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING**

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ sinus congestion and pressure ■ cough due to minor throat and bronchial irritation ■ minor aches and pains ■ headache ■ fever ■ sore throat ■ reduces swelling of nasal passages ■ temporarily restores freer breathing through the nose ■ promotes nasal and/or sinus drainage ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if you have ■ liver disease ■ heart disease ■ thyroid disease ■ high blood pressure ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ diabetes ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ a sodium-restricted diet



3184

DAY & NIGHT PACK

MCC 8168-1709-01



Compare to the active ingredients in **Theraflu® Multi-Symptom Severe Cold & Flu** and **Nighttime Severe Cold & Flu**

MULTI-SYMPTOM Severe Cold DM

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
DOXYLMEPHETHAN HBr / COUGH SUPPRESSANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

- Reduces nasal congestion, sore throat, pain, cough, headache, body aches, fever, stuffy nose & sneezing

6 PACKETS Green Tea & Honey Lemon Flavors

NIGHTTIME Severe Cold & Flu

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
DOXYLMEPHETHAN HCl / ANTIHISTAMINE / COUGH SUPPRESSANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

- Reduces nasal congestion, sore throat, pain, cough, headache, body aches, fever, stuffy nose & sneezing

6 PACKETS Honey Lemon & White Tea Flavors

TOTAL 12 PACKETS

DO NOT TAKE BOTH PRODUCTS AT THE SAME TIME OR TAKE MORE THAN 5 DOSES IN TOTAL IN ANY 24 HOUR PERIOD

DO NOT TAKE A DOSE OF THE NIGHTTIME PRODUCT SOONER THAN 4 HOURS AFTER THE LAST DOSE OF MULTI-SYMPTOM PRODUCT UNLESS DIRECTED BY YOUR DOCTOR

Daytime Multi-Symptom Severe Cold

Drug Facts

Active Ingredients (in each packet):

- Acetaminophen 500 mg Pain reliever/fever reducer
- Doxylamine HCl 20 mg Cough suppressant
- Phenylephrine HCl 10 mg Nasal decongestant

Other Ingredients: Includes inactive ingredients such as citric acid, sodium citrate, and others.

Warnings: Do not take if you have liver disease, kidney disease, or are taking other drugs containing acetaminophen.

Directions: Take 1 packet every 4 hours while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

Severe Cold & Flu NIGHTTIME

Walgreens

Drug Facts (continued)

Other Information: Contains 25 mg of phenylephrine. Contains 22 mg of doxylamine HCl. Contains 10 mg of acetaminophen.

Directions: Take 1 packet every 4 hours while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

Questions or comments? 1-800-746-6200

Drug Facts (continued)

Other Information: Contains 25 mg of phenylephrine. Contains 22 mg of doxylamine HCl. Contains 10 mg of acetaminophen.

Directions: Take 1 packet every 4 hours while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

Questions or comments? 1-800-746-6200

DO NOT TAKE BOTH PRODUCTS AT THE SAME TIME OR TAKE MORE THAN 5 DOSES IN TOTAL IN ANY 24 HOUR PERIOD

DO NOT TAKE A DOSE OF THE NIGHTTIME PRODUCT SOONER THAN 4 HOURS AFTER THE LAST DOSE OF MULTI-SYMPTOM PRODUCT UNLESS DIRECTED BY YOUR DOCTOR

Daytime Multi-Symptom Severe Cold

Drug Facts

Active Ingredients (in each packet):

- Acetaminophen 500 mg Pain reliever/fever reducer
- Doxylamine HCl 20 mg Cough suppressant
- Phenylephrine HCl 10 mg Nasal decongestant

Other Ingredients: Includes inactive ingredients such as citric acid, sodium citrate, and others.

Warnings: Do not take if you have liver disease, kidney disease, or are taking other drugs containing acetaminophen.

Directions: Take 1 packet every 4 hours while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

DO NOT TAKE BOTH PRODUCTS AT THE SAME TIME OR TAKE MORE THAN 5 DOSES IN TOTAL IN ANY 24 HOUR PERIOD

DO NOT TAKE A DOSE OF THE NIGHTTIME PRODUCT SOONER THAN 4 HOURS AFTER THE LAST DOSE OF MULTI-SYMPTOM PRODUCT UNLESS DIRECTED BY YOUR DOCTOR

Daytime Multi-Symptom Severe Cold

Drug Facts

Active Ingredients (in each packet):

- Acetaminophen 500 mg Pain reliever/fever reducer
- Doxylamine HCl 20 mg Cough suppressant
- Phenylephrine HCl 10 mg Nasal decongestant

Other Ingredients: Includes inactive ingredients such as citric acid, sodium citrate, and others.

Warnings: Do not take if you have liver disease, kidney disease, or are taking other drugs containing acetaminophen.

Directions: Take 1 packet every 4 hours while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

DO NOT TAKE BOTH PRODUCTS AT THE SAME TIME OR TAKE MORE THAN 5 DOSES IN TOTAL IN ANY 24 HOUR PERIOD

DO NOT TAKE A DOSE OF THE NIGHTTIME PRODUCT SOONER THAN 4 HOURS AFTER THE LAST DOSE OF MULTI-SYMPTOM PRODUCT UNLESS DIRECTED BY YOUR DOCTOR

Daytime Multi-Symptom Severe Cold

Drug Facts

Active Ingredients (in each packet):

- Acetaminophen 500 mg Pain reliever/fever reducer
- Doxylamine HCl 20 mg Cough suppressant
- Phenylephrine HCl 10 mg Nasal decongestant

Other Ingredients: Includes inactive ingredients such as citric acid, sodium citrate, and others.

Warnings: Do not take if you have liver disease, kidney disease, or are taking other drugs containing acetaminophen.

Directions: Take 1 packet every 4 hours while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

DO NOT TAKE BOTH PRODUCTS AT THE SAME TIME OR TAKE MORE THAN 5 DOSES IN TOTAL IN ANY 24 HOUR PERIOD

DO NOT TAKE A DOSE OF THE NIGHTTIME PRODUCT SOONER THAN 4 HOURS AFTER THE LAST DOSE OF MULTI-SYMPTOM PRODUCT UNLESS DIRECTED BY YOUR DOCTOR

Daytime Multi-Symptom Severe Cold

Drug Facts

Active Ingredients (in each packet):

- Acetaminophen 500 mg Pain reliever/fever reducer
- Doxylamine HCl 20 mg Cough suppressant
- Phenylephrine HCl 10 mg Nasal decongestant

Other Ingredients: Includes inactive ingredients such as citric acid, sodium citrate, and others.

Warnings: Do not take if you have liver disease, kidney disease, or are taking other drugs containing acetaminophen.

Directions: Take 1 packet every 4 hours while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015


100% SATISFACTION GUARANTEED

walgreens.com


©2023 Walgreen Co. MADE IN MEXICO

CODE AREA

1895 04 C3



Compare to active ingredients in
Maximum Strength Mucinex®
Fast-Max® Severe Congestion & Cough™




Severe Congestion & Cough

DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
GUAIFENESIN / EXPECTORANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

Maximum Strength
Multi-Symptom

- Controls cough
- Relieves nasal & chest congestion
- Thins & loosens mucus




20
COATED
CAPLETS



ACTUAL SIZE

Do not print

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING

BOX
MULTI-LAYER
PAPER

50844 ORG042384809
FAST-MAX® Severe Congestion & Cough™
owner of the registered trademark Mucinex®
This product is not manufactured or distributed by B&B Health (US) LLC.
We invite you to compare to national brands.
Our pharmacist recommends the Walgreens brand.

B-2201-648-09-HR
ORG042384809

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredients (in each caplet)

Dextromethorphan HBr 10 mg Cough suppressant

Guaifenesin 200 mg Expectorant

Phenylephrine HCl 5 mg Nasal decongestant

Uses ■ temporarily relieves:

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the intensity of coughing
- nasal congestion due to a cold
- the impulse to cough to help you get to sleep
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- high blood pressure
- heart disease
- diabetes
- thyroid disease
- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than directed

- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years and over: take 2 caplets every 4 hours
- children under 12 years: do not use

Other information ■ each caplet contains: sodium 3 mg

■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

■ see end flap for expiration date and lot number

Purpose

Uses

Warnings

Ask a doctor before use if you have

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Other information

Drug Facts (continued)

Inactive ingredients corn starch, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments? 1-800-426-9391

W3ORG0523-F
REV0523

ITEM 766690 W00000-0000-0
DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
©2023 Walgreen Co.



F-004-45 ORG

PARENTS:

Learn more about our products at
www.DrugFactsNow.org

PEEL BACK TAB TO READ COMPLETE
DRUG FACTS AND INFORMATION

50844 ORG042300445
B-004-45 ORG

DISTRIBUTED BY:
WALGREEN CO.
DEERFIELD, IL 60015
100% SATISFACTION
GUARANTEED
walgreens.com
©2023 Walgreen Co.

ITEM 769446 W00000-0000-0



W3ORG0923-F

*Our pharmacists recommend the Walgreens brand.
*We invite you to compare to national brands.

†This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Mucinex® FAST-MAX® Severe Congestion & Cough.

Drug Facts

TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Active ingredients (in each 20 mL) Purpose

Dextromethorphan HBr 20 mg Cough suppressant
Guaifenesin 400 mg Expectorant
Phenylephrine HCl 10 mg Nasal decongestant

Uses ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
■ temporarily relieves:
■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
■ the intensity of coughing
■ the impulse to cough to help you get to sleep
■ nasal congestion due to a cold

No Print / No Varnish Area
Lot # and Exp. Info

Drug Facts (continued)**Warnings****Do not use**

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

■ high blood pressure ■ thyroid disease
■ difficulty in urination due to enlargement of the prostate gland ■ heart disease ■ diabetes
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ cough that occurs with too much phlegm (mucus)

When using this product do not exceed recommended dosage.**Stop use and ask a doctor if**

■ nervousness, dizziness, or sleeplessness occur
■ symptoms do not get better within 7 days or occur with fever
■ cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)**Directions**

■ do not take more than directed
■ do not take more than 6 doses in any 24-hour period
■ mL = milliliter
■ only use the dose cup provided
■ adults and children 12 years and over: 20 mL in dosing cup provided every 4 hours
■ children under 12 years: do not use

Other information

■ each 20 mL contains: sodium 9 mg
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ use by expiration date on package

Inactive ingredients

anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sorbitol, sucralose, xanthan gum

Questions or comments?

1-800-426-9391

Discard Seal,
Empty &
Replace Cap



PLASTIC
BOTTLE



PLASTIC
CUP

how2recycle.info

PARENTS: **WARNING:** This product contains acetaminophen. Do not give to children under 18 years of age without a doctor's advice. See important information about acetaminophen on page 2.

Walgreens

Compare to the active ingredients in Mucinex® Easy-Max™ Serious Congestion & Cough or Robitussin® Cold & Flu

Severe Congestion & Cough

DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
GLAFENESIN / EXPECTORANT
PHENTYLEPRINE HCl NASAL DECONGESTANT

Maximum Strength

- Controls cough
- Relieves nasal & chest congestion
- Thins & loosens mucus
- 12 years & older

NIGHTTIME Cold & Flu

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
TRIPROLOLINE HCl / ANTIHISTAMINE

Multi-Symptom

- Relieves cough, fever, sore throat, runny nose & sneezing
- 12 years & older

2 - 6 FL OZ (177 mL) BOTTLES / TOTAL 12 FL OZ (355 mL)

No print/No varnish area
Lot no/Exp date

Do not print
Lot & Exp

DISTRIBUTED BY: WALGREENS CO., 200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
Walgreens.com/2022WalgreenCo.

TAMPER EVIDENT: DO NOT USE IF PACKAGING IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

50844 ORG092252708

NDC 0363-5027-08



Compare to the active ingredients in Tylenol® Sinus Severe™

DAYTIME • NON-DROWSY

Severe Sinus

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
GUAIFENESIN / EXPECTORANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

Severe

• Relieves sinus headache, nasal congestion, sinus pressure, chest congestion & mucus

24
CAPLETS



ACTUAL SIZE

Drug Facts (continued)

Other Information ■ each caplet contains sodium 3 mg
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ see end flap for expiration date and lot number

Drug Facts (continued)

Inactive ingredients corn starch, croscollon, fluor magnesium stearate, maleic acid, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pyridone, silicon dioxide, sodium stearoyl glycolate, stearic acid, sucralose, etc., titanium dioxide

Questions or comments? 1-800-426-9391

Drug Facts (continued)

Directions

- do not take more than directed
- adults and children 12 years and over
- take 2 caplets every 4 hours
- swallow whole- do not crush, chew or dissolve
- do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Warnings

- liver disease ■ high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if:

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

W3ORG0722-F
REV1222

ITEM 510073 W00000-0000-0



119171103723

B-2201-527C-08SSVR
ORG092252708



NDC 0363-4666-08

DAYTIME • NON-DROWSY

Sinus & Headache

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
PHENYLEPHRINE HCl / NASAL DECONGESTANT

- Relieves hay fever symptoms, minor aches & pains, headache, nasal congestion, sinus congestion & pressure
- Pseudoephedrine free

24
CAPLETS

ACTUAL SIZE

Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active Ingredients (in each caplet)

Acetaminophen 325 mg Pain reliever/fever reducer
Phenylephrine HCl 5 mg Nasal decongestant

Purpose

Uses

- temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold:
- minor aches and pains ■ headache ■ nasal congestion
- sinus congestion and pressure

Drug Facts (continued)

- helps decongest sinus openings and passages
- promotes sinus drainage
- helps clear nasal passages
- temporarily reduces fever

Warnings

Live warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 1,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 5 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ rash ■ skin reddening
■ blisters ■ skin reaction occurs, stop use and seek medical help right away.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- difficulty in urination due to enlarged prostate of the prostate gland
- liver disease ■ heart disease ■ hay fever ■ high blood pressure

Drug Facts (continued)

Inactive ingredients: corn starch, croscarmellose, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose, polyethylene glycol, polyvinyl alcohol, povidone, simethicone, sodium starch glycolate, stearic acid, sucrose, etc., titanium dioxide

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- these could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- adults and children 12 years and over
- take 2 caplets every 4 hours
- swallow whole – do not crush, chew, or dissolve
- do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

■ **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTERS ARE TORN OR BROKEN**
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ see end flap for expiration date and lot number

Drug Facts (continued)

(Questions or comments? 1-800-226-9391)

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
©2022 Walgreen Co.
50844 ORG072146608

B-2201-466C-08SHVR
ORG072146608

W3ORG0722-F

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
This product is not manufactured or distributed by Johnson & Johnson trademark Tylenol® SINUS + HEADACHE.

Do not print
Lot & Exp

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTERS ARE TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

NDC 0363-5580-08

DAYTIME • NON-DROWSY

Sinus & Headache

**ACETAMINOPHEN /
PAIN RELIEVER / FEVER REDUCER
PHENYLEPHRINE HCl / NASAL DECONGESTANT**

- Relieves headache, nasal congestion & sinus pressure

Compare to Tylenol® Sinus + Headache Daytime active ingredients!

Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredients
(in each gelcap)
Acetaminophen 325 mg Pain reliever/fever reducer
Phenylephrine HCl 5 mg Nasal decongestant

Purpose
Relieves sinus pressure, nasal congestion, and the common cold; headache associated with fever or other respiratory symptoms

Uses
temporarily relieves these symptoms associated with fever or other respiratory symptoms: ■ headache ■ nasal congestion ■ sinus congestion and pressure

Drug Facts (continued)

Drug Facts (continued)

■ difficulty in urination due to enlargement of the prostate gland ■ high blood pressure

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occur ■ pain or nasal congestion gets worse or lasts more than 7 days ■ new symptoms occur ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present

These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
■ do not take more than directed ■ adults and children 12 years and over ■ take 2 gelcaps every 4 hours ■ do not take more than 10 gelcaps in 24 hours ■ children under 12 years: ask a doctor

Other information
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN ■ avoid high humidity ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ see end flap for expiration date and lot number

Drug Facts (continued)

Inactive ingredients
croscarmellose sodium, croscarmellose, iron oxide black, iron oxide red #40, FD&C yellow #6, gelatin, hydroxypropyl cellulose, hypromellose, iron oxide red, iron oxide yellow, magnesium stearate.

Drug Facts (continued)

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
■ liver disease ■ heart disease ■ thyroid disease ■ diabetes

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
■ liver disease ■ heart disease ■ thyroid disease ■ diabetes

Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey
This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® Sinus + Headache Daytime.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015

Walgreens

**100% SATISFACTION
GUARANTEED**

walgreens.com ©2017 Walgreen Co.

50844 ORG021655808

B-2201-558-08-H
ORG021655808




TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING


No Print Area
Lot no. and expiration date

Drug Facts (continued)

Questions or comments? 1-800-426-9391

ORG0917-F





W00R61221-F
REV0122

DISTRIBUTED BY: WALGREEN CO.
200 WILLIOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
Walgreens.com
©2021 Walgreen Co.

ITEM 357646 W00000-0000-0
11917116860 9

B-2201-628-09-HR
OR6041962809

Walgreens
Compare to the active ingredients
in Maximum Strength Mucinex®
SINUS-MAX® Pressure & Pain

Sinus Pressure & Pain

ACETAMINOPHEN 325 mg / PAIN RELIEVER
GUAIFENESIN 200 mg / EXPECTORANT
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

Maximum Strength

• Relieves sinus pressure & congestion, headache
and thins & loosens mucus

20
CAPLETS

ACTUAL SIZE

NDC 0363-69-028-09

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredients (in each caplet)
Acetaminophen 325 mg Pain reliever
Guaifenesin 200 mg Expectorant
Phenylephrine HCl 5 mg Nasal decongestant

Purpose

• temporarily relieves:
■ sinus congestion and pressure
■ nasal congestion
■ headache

Uses

■ temporarily relieves:
■ sinus congestion and pressure
■ nasal congestion
■ headache

Drug Facts (continued)

■ temporarily relieves:
■ sinus congestion and pressure
■ nasal congestion
■ headache

Drug Facts (continued)

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ blisters ■ rash ■ skin redness
If a skin reaction occurs, stop use and seek medical help right away.

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
■ heart disease ■ diabetes ■ thyroid disease
■ difficulty in urination due to enlargement of the prostate gland
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema ■ liver disease
■ cough that occurs with too much phlegm (mucus)

Inactive ingredients corn starch, croscopolone, D.C. yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Drug Facts (continued)

Other information ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ see end flap for expiration date and lot number

Directions
■ do not take more than directed
■ adults and children 12 years and over: take 2 caplets every 4 hours
■ children under 12 years: do not use

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

When using this product do not exceed recommended dosage.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Drug Facts (continued)

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occur
■ fever gets worse or lasts more than 3 days
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
■ redness or swelling is present
■ new symptoms occur

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
■ heart disease ■ diabetes ■ thyroid disease
■ difficulty in urination due to enlargement of the prostate gland
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema ■ liver disease
■ cough that occurs with too much phlegm (mucus)

Inactive ingredients corn starch, croscopolone, D.C. yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Drug Facts (continued)

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occur
■ fever gets worse or lasts more than 3 days
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
■ redness or swelling is present
■ new symptoms occur

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
■ heart disease ■ diabetes ■ thyroid disease
■ difficulty in urination due to enlargement of the prostate gland
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema ■ liver disease
■ cough that occurs with too much phlegm (mucus)

Inactive ingredients corn starch, croscopolone, D.C. yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Drug Facts (continued)

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occur
■ fever gets worse or lasts more than 3 days
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
■ redness or swelling is present
■ new symptoms occur

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
■ heart disease ■ diabetes ■ thyroid disease
■ difficulty in urination due to enlargement of the prostate gland
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema ■ liver disease
■ cough that occurs with too much phlegm (mucus)

Inactive ingredients corn starch, croscopolone, D.C. yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Drug Facts (continued)

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occur
■ fever gets worse or lasts more than 3 days
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
■ redness or swelling is present
■ new symptoms occur

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
■ heart disease ■ diabetes ■ thyroid disease
■ difficulty in urination due to enlargement of the prostate gland
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema ■ liver disease
■ cough that occurs with too much phlegm (mucus)

Inactive ingredients corn starch, croscopolone, D.C. yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Drug Facts (continued)

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occur
■ fever gets worse or lasts more than 3 days
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
■ redness or swelling is present
■ new symptoms occur

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
■ heart disease ■ diabetes ■ thyroid disease
■ difficulty in urination due to enlargement of the prostate gland
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema ■ liver disease
■ cough that occurs with too much phlegm (mucus)

Inactive ingredients corn starch, croscopolone, D.C. yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Drug Facts (continued)

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occur
■ fever gets worse or lasts more than 3 days
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
■ redness or swelling is present
■ new symptoms occur

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
■ heart disease ■ diabetes ■ thyroid disease
■ difficulty in urination due to enlargement of the prostate gland
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema ■ liver disease
■ cough that occurs with too much phlegm (mucus)

Inactive ingredients corn starch, croscopolone, D.C. yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

No Print / No Varnish
Lot no. & Exp. date

No print/No varnish area

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by Walgreens (US) LLC, owner of the registered trademark, Maximum Strength Mucinex®. SINUS-MAX Day & Night Maximum Strength Mucinex®. 50844 ORG051761569409

W3ORG0423-F

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2023 Walgreen Co.

ITEM 886729 W00000-0000-0

3 11917 19527 8

PEEL HERE FOR MORE DRUG FACTS

Drug Facts
KEEP OUT OF REACH OF CHILDREN
COMPLETE PRODUCT INFORMATION

Active ingredients (in each caplet)
Acetaminophen 325 mg
Pain reliever

Active ingredients (in each caplet)
SINUS-MAX Day & Night
Acetaminophen 325 mg
Pain reliever

Other ingredients (in each caplet)
Diphenhydramine HCl 12.5 mg
Antihistamine/cough suppressant

Other ingredients (in each caplet)
SINUS-MAX Day & Night
Diphenhydramine HCl 12.5 mg
Antihistamine/cough suppressant

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Directions
Take 1 caplet 4 times a day, every 6 hours, with or without food. Do not take more than 12 caplets in any 24-hour period. Adults and children 12 years and over: take 2 caplets every 4 hours. Children under 12 years: do not use.

Other information
TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Inactive ingredients (Daytime only)
corn starch, croscarmellose sodium, croscopolone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, iron oxide yellow, magnesium stearate, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Inactive ingredients (Nighttime only)
corn starch, croscarmellose sodium, croscopolone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, iron oxide yellow, magnesium stearate, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments? 1-800-426-9391

Walgreens Pharmacist Recommended. Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2023 Walgreen Co.

50844
ORG051761569409
R-2201-615694-09HRV-DF

W3ORG0423-F

DAY & NIGHT PACK NDC 0363-6156-09

Walgreens
Compare to the active ingredients in Maximum Strength Mucinex® SINUS-MAX® Day & Night™

DAYTIME
Sinus Pressure & Pain
ACETAMINOPHEN
PAIN RELIEVER
GUAIFENESIN
EXPECTORANT
PHENYLEPHRINE HCl
NASAL DECONGESTANT
Maximum Strength

NIGHTTIME
Sinus Pressure & Pain
ACETAMINOPHEN
PAIN RELIEVER
DIPHENHYDRAMINE HCl
ANTIHISTAMINE
COUGH SUPPRESSANT
PHENYLEPHRINE HCl
NASAL DECONGESTANT
Maximum Strength

12 CAPLETS ACTUAL SIZE 8 CAPLETS ACTUAL SIZE
TOTAL 20 CAPLETS

Do not take the Daytime and Nighttime caplets at the same time.

B-2201-615694-09HRV
ORG051761569409

Drug Facts (continued)

■ with any other product containing diphenhydramine, even one used on skin (Nighttime only)

Ask a doctor before use if you have

■ heart disease ■ high blood pressure
■ liver disease ■ thyroid disease ■ diabetes
■ difficulty in urination due to enlargement of the prostate gland
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ cough that occurs with too much phlegm (mucus)
■ a breathing problem such as emphysema or chronic bronchitis (Nighttime only) ■ glaucoma (Nighttime only)

Ask a doctor or pharmacist before use if you are

■ taking the blood thinning drug warfarin
■ taking sedatives or tranquilizers (Nighttime only)

When using this product

■ do not exceed recommended dosage
■ excitability may occur, especially in children (Nighttime only) ■ marked drowsiness may occur (Nighttime only)
■ alcohol, sedatives, and tranquilizers may increase drowsiness (Nighttime only)

Drug Facts (continued)

■ be careful when driving a motor vehicle or operating machinery (Nighttime only)

■ avoid alcoholic beverages (Nighttime only)

Stop use and ask a doctor if

■ nervousness, dizziness, or sleeplessness occur
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ new symptoms occur
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

If taking NIGHTTIME and DAYTIME products, carefully read each section to ensure correct dosing. Do not take DAY & NIGHT at the same time.

Drug Facts (continued)

Directions

■ do not use more than directed
■ do not take more than 12 caplets in any 24-hour period
■ adults and children 12 years and over: take 2 caplets every 4 hours
■ children under 12 years: do not use

Other information

■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ see end flap for expiration date and lot number

Inactive ingredients (Daytime only)
corn starch, croscopolone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Inactive ingredients (Nighttime only)
corn starch, croscarmellose sodium, croscopolone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, iron oxide yellow, magnesium stearate, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium bicarbonate, stearic acid, talc, titanium dioxide

Questions or comments? 1-800-426-9391

Walgreens Pharmacist Recommended. Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2023 Walgreen Co.

50844
ORG051761569409
R-2201-615694-09HRV-DF

W3ORG0423-F

DAY & NIGHT PACK NDC 0363-6156-10


Compare to the active ingredients
in Maximum Strength Mucinex[®]
SINUS-MAX[®] Day & Night[®]



DAYTIME

Sinus Pressure & Pain

ACETAMINOPHEN
PAIN RELIEVER
GUAIFENESIN
EXPECTORANT
PHENYLEPHRINE HCl
NASAL DECONGESTANT

Maximum Strength


24 CAPLETS ACTUAL SIZE

NIGHTTIME

Sinus Pressure & Pain

ACETAMINOPHEN
PAIN RELIEVER
DIPHENHYDRAMINE HCl
ANTIHISTAMINE
COUGH SUPPRESSANT
PHENYLEPHRINE HCl
NASAL DECONGESTANT

Maximum Strength


16 CAPLETS ACTUAL SIZE

TOTAL 40 CAPLETS

Do not take the Daytime and Nighttime caplets at the same time.

B-2201-61560410HRY
CR0251761560410

PAIN EXHAUSTIVE

Drug Facts (continued)

- with any other product containing diphenhydramine, even one used on skin (*Nighttime only*)

Ask a doctor before use if you have

- heart disease
- high blood pressure
- liver disease
- thyroid disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis (*Nighttime only*)
- glaucoma (*Nighttime only*)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (*Nighttime only*)

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children (*Nighttime only*)
- marked drowsiness may occur (*Nighttime only*)
- alcohol, sedatives, and tranquilizers may increase drowsiness (*Nighttime only*)

PAIN EXHAUSTIVE

Drug Facts (continued)

- be careful when driving a motor vehicle or operating machinery (*Nighttime only*)
- avoid alcoholic beverages (*Nighttime only*)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that
- could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

If taking NIGHTTIME and DAYTIME products, carefully read each section to ensure correct dosing. Do not take DAY & NIGHT at the same time.

PAIN EXHAUSTIVE

Drug Facts (continued)

Directions

- do not use more than directed
- do not take more than 12 capsules in any 24-hour period
- adults and children 12 years and over: take 2 capsules every 4 hours
- children under 12 years: do not use

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients (Daytime only)

corn starch, croscopolone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

PAIN EXHAUSTIVE

Drug Facts (continued)

Inactive ingredients (Nighttime only)

corn starch, croscopolone sodium, croscopolone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, iron oxide yellow, magnesium stearate, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium bicarbonate, stearic acid, talc, titanium dioxide


Questions or comments? 1-800-426-9391

Walgreens Pharmacist Recommended.

Our pharmacists recommend the Walgreens brand. We invite you to compare us to national brands.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD.,
DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
(800) 521-7616
R-2201-61569410HRY-DF

ITEM 355984
W00000-0000-0



W3RG0G0423-F

Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredients (in each caplet)

Acetaminophen 325 mg.....Pain reliever
 Dextromethorphan HBr 10 mg.....Cough suppressant
 Guaifenesin 200 mg.....Expectorant
 Phenylephrine HCl 5 mg.....Nasal decongestant

Purpose

Drug Facts (continued)

Uses ■ temporarily relieves: ■ nasal congestion ■ headache
 ■ sinus congestion and pressure ■ minor aches and pains
 ■ cough due to the common cold
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive ■ temporarily promotes nasal and/or sinus drainage

Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ blisters ■ rash ■ skin reddening
 If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- heart disease ■ diabetes ■ thyroid disease
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ liver disease ■ high blood pressure
- cough that occurs with too much phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present ■ new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years and over: take 2 caplets every 4 hours
- children under 12 years: do not use

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- see end flap for expiration date and lot number

Drug Facts (continued)

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients corn starch, croscopolone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl

Drug Facts (continued)

alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments? 1-800-426-9391

how2recycle.info

PARENTS:

Learn about how medicine affects your child's health at www.StopMedicineAbuse.org

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

DISTRIBUTED BY: WALGREENS CO.
 200 WILMOT RD., DEERFIELD, IL 60015
 100% SATISFACTION GUARANTEED
 walgreens.com ©2022 Walgreen Co.

W3ORG0422-F REV1022

Walgreens

Compare to the active ingredients
 in Maximum Strength Mucinex®
 SINUS-MAX® Pressure, Pain & Cough

NDC 0363-6171-09

Sinus Pressure, Pain & Cough

ACETAMINOPHEN / PAIN RELIEVER
 DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
 GUAIFENESIN / EXPECTORANT
 PHENYLEPHRINE HCl / NASAL DECONGESTANT

Maximum Strength

- Relieves headache, sinus pressure & congestion
- Controls cough
- Thins & loosens mucus

20

CAPLETS

ACTUAL SIZE



Drug Facts

Active ingredients (in each 20 mL) **Purposes**

Acetaminophen, USP 650 mg Pain reliever/fever reducer

Dextromethorphan HBr, USP 20 mg Cough suppressant

Guaifenesin, USP 400 mg Expectorant

Phenylephrine HCl, USP 10 mg Nasal decongestant

Uses

- temporarily relieves these symptoms occurring with a cold or flu:
 - cough due to minor throat and bronchial irritation
 - nasal congestion
 - chest congestion and pressure
 - minor aches and pains
 - sore throat
 - headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- hives
- rash

If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before using this product.
- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than 5 doses in any 24-hour period
- do not exceed recommended dosage. Taking more than the recommended dose (overdose) may cause serious liver damage.
- measure only with dosing cup provided
- keep dosing cup with product
- 1 mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

Other information

- each 20 mL contains: sodium 14 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients anhydrous citric acid, benzyl alcohol, edetate disodium, FD&C red #40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, xanthane gum, xanthan gum

Questions or comments? 1-800-719-9280

*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
**This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare Holdings (US) LLC, owner of the registered trademark Robitussin®.

Empty & Recycle Cup
Check Locally
PAPER BOX PLASTIC BOTTLE PLASTIC CUP
*not recycled in all communities

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Walgreens

ADULT • NON-DROWSY

Tussin CF Max

MULTI-SYMPTOM
COUGH, COLD & FLU

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

NDC 0363-2801-34

Compare to the active ingredients in Robitussin® Severe Multi-Symptom Cough Cold + Flu*

ADULT NON-DROWSY

Tussin CF Max

MULTI-SYMPTOM
COUGH, COLD & FLU

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
GUAIFENESIN / EXPECTORANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

Maximum Strength

- Relieves cough, sore throat, body aches, fever, nasal congestion & chest congestion
- Ages 12 & older

8 FL OZ (237 mL) Raspberry flavor

DISTRIBUTED BY: WALGREEN CO.
200 WILLMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
©2022 Walgreen Co.

ITEM 838110 W10106-0323-L

28S34 94 C1

Walgreens

ADULT • NON-DROWSY

Tussin CF Max

MULTI-SYMPTOM
COUGH, COLD & FLU

NDC 0363-2801-34

Compare to the active ingredients in Robitussin® Severe Multi-Symptom Cough Cold + Flu*

ADULT NON-DROWSY

Tussin CF Max

MULTI-SYMPTOM
COUGH, COLD & FLU

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
GUAIFENESIN / EXPECTORANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

Maximum Strength

- Relieves cough, sore throat, body aches, fever, nasal congestion & chest congestion
- Ages 12 & older

8 FL OZ (237 mL) Raspberry flavor

DAY & NIGHT PACK

Walgreens

Wal-Flu® Severe

COLD

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

6 PACKETS

GREEN TEA & HONEY LEMON FLAVORS

Walgreens

Wal-Flu® Severe

COLD & COUGH

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
DIPHENHYDRAMINE HCl / ANTIHISTAMINE / COUGH SUPPRESSANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

6 PACKETS

HONEY LEMON INFUSED WITH WHITE TEA FLAVORS

TOTAL 12 PACKETS

DO NOT TAKE BOTH PRODUCTS AT THE SAME TIME OR TAKE MORE THAN 5 DOSES IN TOTAL IN ANY 24 HOUR PERIOD

DO NOT TAKE A DOSE OF THE NIGHTTIME PRODUCT SOONER THAN 4 HOURS AFTER THE LAST DOSE OF MULTI-SYMPTOM PRODUCT UNLESS DIRECTED BY YOUR DOCTOR

Daytime Multi-Symptom Severe Cold

Drug Facts

Active Ingredients (in each packet)

Acetaminophen 500 mg... **Purpose:** This helps relieve pain and fever.

Dextromethorphan HCl 15 mg... **Purpose:** Cough suppressant.

Phenylephrine HCl 10 mg... **Purpose:** Nasal decongestant.

Uses: It temporarily relieves these symptoms due to a cold: fever, aches and pain, sore throat, runny nose, sneezing, headache, and congestion.

Warnings: Liver warning: This product contains acetaminophen. Serious liver damage may occur if you take: more than 4,000 mg of acetaminophen in 24 hours; 3 or more alcoholic drinks a day while using this product; or with other drugs containing acetaminophen.

Directions: Use as directed. Do not take more than 5 doses in 24 hours.

Walgreens

Wal-Flu® Severe

NIGHTTIME

COLD & COUGH

Drug Facts (continued)

Other Information: Do not use if you are taking a prescription medicine for depression, seizure, or heart disease.

Directions (continued): Do not take more than 5 doses in 24 hours.

Walgreens

Wal-Flu® Severe

NIGHTTIME

COLD & COUGH

Drug Facts (continued)

Other Information: Do not use if you are taking a prescription medicine for depression, seizure, or heart disease.

Directions (continued): Do not take more than 5 doses in 24 hours.

Walgreens

Wal-Flu® Severe

NIGHTTIME

COLD & COUGH

Drug Facts (continued)

Other Information: Do not use if you are taking a prescription medicine for depression, seizure, or heart disease.

Directions (continued): Do not take more than 5 doses in 24 hours.

DISTRIBUTED BY: WALGREEN CO.

201 WILSON ST., GAITHERSBURG, MD 20878

100% SATISFACTION GUARANTEED

Walgreens

MADE IN MEXICO

ITEM 443265 W10105-1214-L

119177-21208



Compare to Theraflu® Multi-Symptom Severe Cold active ingredients!

NDC 0983-8219-01

MULTI-SYMPTOM Wal-Flu® Severe COLD

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

- Relieves nasal congestion, sore throat pain, cough, headache, body ache & fever



6 PACKETS

GREEN TEA & HONEY LEMON FLAVORS

PHARMACIST



Trusted since 1901™

RECOMMENDED

Health expertise you rely on.™

CONVENIENT RESEALING TAB

Walgreens

DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED

Drug Facts

Active ingredients (in each packet)

Acetaminophen 500 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold:
 - minor aches and pains
 - minor sore throat pain
 - headache
 - nasal and sinus congestion
 - cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours.

Severe throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- hives
- itching

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not use more than directed (see overdose warning)
- take every 4 hours, while symptoms persist. Do not take more than 6 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and over	one packet
children under 12 years of age	do not use

Other information

- each packet contains: potassium 10 mg and sodium 25 mg
- phenylephrine contains phenylalanine
- 22 mg per packet
- store at 20-25°C (68-77°F). Protect product from heat and moisture.

Inactive ingredients acetaminophen, anhydrous citric acid, aspartame, colloidal silicon dioxide, DSC yellow #10, FD&C blue #1, FD&C red #40, flavors, malachite green, pregelatinized starch, sodium citrate, sucrose, tribasic calcium phosphate

Questions or comments?
1-800-719-8280

Gluten Free
Walgreens Pharmacist Survey
This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Theraflu® Multi-Symptom Severe Cold.





ORG0518-F
REV-1019

OPEN OTHER END

DISTRIBUTED BY: WALGREEN CO.
300 WILMOT RD., DEERFIELD, IL 60015

Walgreens

100% SATISFACTION GUARANTEED

walgreens.com ©2018 walgreen co.
MADE IN MEXICO

ITEM 038993 W10106-1219-L



3 11917 21201 2

8Y491 94 C2



CONVENIENT RECLOSED TAB



Wal-Fin
NIGHTIME
Severe Cold & Cough

DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED

Drug Facts

Active ingredients (in each packet)

Acetaminophen 650 mg.....	Pain reliever/fever reducer
Diphenhydramine HCl 25 mg.....	Antihistamine/cough suppressant
Phenylephrine HCl 10 mg.....	Nasal decongestant

Purposes

Uses ■ temporarily relieves these symptoms due to a cold:

- minor aches and pains
- minor sore throat pain
- headache
- nasal and sinus congestion
- runny nose
- sneezing
- itchy nose or throat
- itchy, watery eyes due to hay fever
- cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ in a child under 12 years of age

- If you have ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ liver disease ■ heart disease

■ high blood pressure ■ thyroid disease ■ diabetes ■ glaucoma

- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers

■ taking the blood thinning drug warfarin

When using this product ■ do not exceed recommended dosage

- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occurs

- redness or swelling is present
- fever gets worse or lasts more than 3 days
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose

warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not use more than directed (see overdose warning)

- take every 4 hours, while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and over	one packet
children under 12 years of age	do not use

- dissolve contents of one packet into 8 oz. hot water: sip while hot. Consume entire drink within 10-15 minutes.

- If using a microwave, add contents of one packet to 8 oz. of cool water: stir briskly before and after heating. Do not overheat.

Other information

- each packet contains: potassium 10 mg and sodium 25 mg

- phenylketonurics: contains phenylalanine 13 mg per packet

- store at 20-25°C (68-77°F). Protect product from heat and moisture.

Inactive ingredients

acesulfame potassium, anhydrous citric acid, aspartame, colloidal silicon dioxide, D&C yellow #10, FD&C blue #1, FD&C red #40, flavors, maltodextrin, pregelatinized starch, sodium citrate, sucrose, tribasic calcium phosphate

Questions or comments?

1-800-719-9260

Gluten Free

OPEN OTHER END

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015

Walgreens

ITEM 912518 W10106-0217-L





3207

01616144

Wang says Rhinoceros Harem mated. "This product is not manufactured odor-free but only when you Rhinoceros for day."

Drug facts (cont'd in use)

Other important information

Store & control conditions: Store at controlled room temperature 15° to 30° (59° to 86°F).

Inactive ingredients Sodium chloride and Na₂HPO₄ monohydrate, polysorbate 80, polyethylene glycol 400, polyethylene glycol 600, polyethylene glycol 300, polyethylene glycol 200, polyethylene glycol 150, polyethylene glycol 100, polyethylene glycol 75, polyethylene glycol 50, polyethylene glycol 35, polyethylene glycol 25, polyethylene glycol 15, polyethylene glycol 10, polyethylene glycol 5, polyethylene glycol 3, polyethylene glycol 2, polyethylene glycol 1, polyethylene glycol 0.5, polyethylene glycol 0.25, polyethylene glycol 0.125, polyethylene glycol 0.0625, polyethylene glycol 0.03125, polyethylene glycol 0.015625, polyethylene glycol 0.0078125, polyethylene glycol 0.00390625, polyethylene glycol 0.001953125, polyethylene glycol 0.0009765625, polyethylene glycol 0.00048828125, polyethylene glycol 0.000244140625, polyethylene glycol 0.0001220703125, polyethylene glycol 0.00006103515625, polyethylene glycol 0.000030517578125, polyethylene glycol 0.0000152587890625, polyethylene glycol 0.00000762939453125, polyethylene glycol 0.000003814697265625, polyethylene glycol 0.0000019073486328125, polyethylene glycol 0.00000095367431640625, polyethylene glycol 0.000000476837158203125, polyethylene glycol 0.0000002384185791015625, polyethylene glycol 0.00000011920928955078125, polyethylene glycol 0.000000059604644775390625, polyethylene glycol 0.0000000298023223876953125, polyethylene glycol 0.00000001490116119384765625, polyethylene glycol 0.000000007450580596923828125, polyethylene glycol 0.0000000037252902984619140625, polyethylene glycol 0.00000000186264514923095703125, polyethylene glycol 0.000000000931322574615478515625, polyethylene glycol 0.0000000004656612873077392578125, polyethylene glycol 0.00000000023283064365386962890625, polyethylene glycol 0.000000000116415321826934814453125, polyethylene glycol 0.0000000000582076609134674072265625, polyethylene glycol 0.00000000002910383045673370361328125, polyethylene glycol 0.000000000014551915228366851806640625, polyethylene glycol 0.0000000000072759576141834259033203125, polyethylene glycol 0.00000000000363797880709171295166015625, polyethylene glycol 0.000000000001818989403545856475830078125, polyethylene glycol 0.0000000000009094947017729282379150390625, polyethylene glycol 0.00000000000045474735088646141895751953125, polyethylene glycol 0.000000000000227373675443230709478759765625, polyethylene glycol 0.0000000000001136868377216153547393798828125, polyethylene glycol 0.00000000000005684341886080767736968994140625, polyethylene glycol 0.000000000000028421709430403838684844970703125, polyethylene glycol 0.0000000000000142108547152019193424224853515625, polyethylene glycol 0.00000000000000710542735760095967121124267578125, polyethylene glycol 0.000000000000003552713678800479835605621337890625, polyethylene glycol 0.0000000000000017763568394002399178028106689453125, polyethylene glycol 0.00000000000000088817841970011995890140533447265625, polyethylene glycol 0.000000000000000444089209850059979450702667236328125, polyethylene glycol 0.0000000000000002220446049250299897253513336181640625, polyethylene glycol 0.00000000000000011102230246251499486267666680908203125, polyethylene glycol 0.000000000000000055511151231257497431338333340441015625, polyethylene glycol 0.0000000000000000277555756156287487156691666702205078125, polyethylene glycol 0.00000000000000001387778780781437435783458333511025390625, polyethylene glycol 0.000000000000000006938893903907187178917291667555126953125, polyethylene glycol 0.0000000000000000034694469519535935894586458337775634765625, polyethylene glycol 0.00000000000000000173472347597679679472932291688878173828125, polyethylene glycol 0.00000000000000000086736173798834983973646115594443908703125, polyethylene glycol 0.000000000000000000433680868994174919868230577972219543515625, polyethylene glycol 0.0000000000000000002168404344970874599341152889861097717578125, polyethylene glycol 0.000000000000000000108420217248543729967057644493054885890625, polyethylene glycol 0.0000000000000000000542101086242718649835288222465274429453125, polyethylene glycol 0.00000000000000000002710505431213593249176441112326372147265625, polyethylene glycol 0.00000000000000000001355252715606796624588220556163186073828125, polyethylene glycol 0.000000000000000000006776263578033983122941102780815930369140625, polyethylene glycol 0.0000000000000000000033881317890169915614705513904079651845703125, polyethylene glycol 0.0000000000000000000016940658945084957807352756952039825922890625, polyethylene glycol 0.00000000000000000000084703294725424789036763784760199129614453125, polyethylene glycol 0.000000000000000000000423516473627123945183818923800995648072265625, polyethylene glycol 0.0000000000000000000002117582368135619725919094619004978240361328125, polyethylene glycol 0.00000000000000000000010587911840678098629595473095024891201806640625, polyethylene glycol 0.00000000000000000000005293955920339049314797736547512445600903125, polyethylene glycol 0.000000000000000000000026469779601695246573988682737562228004515625, polyethylene glycol 0.0000000000000000000000132348898008476232869943413687811140022578125, polyethylene glycol 0.00000000000000000000000661744490042381164349717068439057200112890625, polyethylene glycol 0.000000000000000000000003308722450211905821748585342195286000564453125, polyethylene glycol 0.0000000000000000000000016543612251059529108742926710976430002822265625, polyethylene glycol 0.00000000000000000000000082718061255297645543721633554882150014111328125, polyethylene glycol 0.000000000000000000000000413590306276488227718608167774410750070556640625, polyethylene glycol 0.000

[illegible][illegible]

Walgreens

DAYTIME • NON-DROWSY

Cold & Flu

Walgreens

DAYTIME • NON-DROWSY

Cold & Flu

ACETAMINOPHEN 325 mg / ACHES / FEVER / SORE THROAT
DEXTROMETHORPHAN HBr 10 mg / COUGH
PHENYLEPHRINE HCl 5 mg / NASAL CONGESTION

MULTI-SYMPTOM LIQUID CAPS

- Relieves aches, fever, sore throat, nasal congestion & cough

16 LIQUID CAPS

Abstract 500

Compare to Vicks® DayQuil®
LiquiCaps® active ingredients††

NDC 0303-1100-15

IT EM 609868 W1000-0000-D



3 11917 20947 0

DISTRIBUTED BY: WALGREEN CO.
 200 WILMOT RD., DEERFIELD, IL 60015
 1-800-368-3673
 100% SATISFACTION GUARANTEED
 walgreens.com ©2019 Walgreen Co.


MADE IN INDIA

MADE IN INDIA

10

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION. THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND TAMPER-EVIDENT PACKAGE. USE ONLY IF BUSTERS ARE INTACT.

LOT NO:
EXP DATE:



ORG0319-F
REV0221

Drug Facts (continued)

Inactive ingredients corn starch, croscopolone, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

(Questions or comments?) 1-800-426-9391

Drug Facts (continued)

Other information

■ **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

■ see end flap for expiration date and lot number

DISTRIBUTED BY: WALGREEN CO.
200 WILLMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2019 Walgreen Co.

ITEM 531245 W00000-0000-0



3 19171 20829 9

Drug Facts (continued)

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

■ cough that occurs with too much phlegm (mucus)

■ Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if:

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- adults and children 12 years and over
- take 2 caplets every 4 hours
- swallow whole - do not crush, chew, or dissolve
- do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Drug Facts (continued)

■ helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ blisters ■ rash ■ skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- heart disease
- liver disease
- diabetes
- thyroid disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland

Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey
This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Sudafed PE® Head Congestion + Mucus

50844 ORG082052708

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

**No Print/No Varnish
Lot and Expiration**

Drug Facts (continued)

Uses

- temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold:
- headache ■ nasal congestion ■ minor aches and pains
- sinus congestion and pressure
- temporarily reduces fever

Drug Facts

Active ingredients (in each caplet)

- Acetaminophen 325 mg.....Pain reliever/fever reducer
- Guaifenesin 200 mg.....Expectorant
- Phenylephrine HCl 5 mg.....Nasal decongestant

Purpose

COMPLETE PRODUCT INFORMATION

B-2201-527C-08HCNH
ORG082052708

Walgreens

NON-DROWSY

Wal-Phed® PE

HEAD CONGESTION + MUCUS

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
GUAIFENESIN / EXPECTORANT / PHENYLEPHRINE HCl / NASAL DECONGESTANT

MULTI-SYMPTOM

- Relieves sinus pressure/congestion, sinus headache & chest congestion
- Pseudoephedrine free

24 CAPLETS



ACTUAL SIZE

Compare to Sudafed PE® Head Congestion + Mucus active ingredients!

NDC 0363-4121-08

Walgreens

Wal-Phed® PE

NASAL DECONGESTANT

BONUS SIZE BUY 18 & GET 9 FREE

NDC 0363-0453-90


Walgreens

Compare to Sudafed PE®
Congestion active ingredient!!

MAXIMUM STRENGTH

- Relieves sinus pressure & congestion
- Pseudoephedrine free

27 TABLETS
(18 + 9 FREE)

 ACTUAL SIZE

Walgreens

100% SATISFACTION
GUARANTEED

200 WILMOT RD., DEERFIELD, IL 60015

walgreens.com ©2018 Walgreen Co.

ITEM 805112 W00000-000-0

11917 20326 3

Walgreens Pharmacist Recommended

Walgreens Sudafed PE® Congestion.

50844 ORG011845372

ORG0918-F

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

DOES NOT CONTAIN GLUTEN

No print area
Lot no/Exp date

Drug Facts

Active ingredient (in each tablet)

Phenylephrine HCl 10 mg Nasal decongestant

Purpose

temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies

temporarily relieves sinus congestion and pressure

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

heart disease ■ diabetes ■ high blood pressure ■ difficulty in urination due to enlargement of the prostate gland

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

■ nervousness, dizziness, or sleeplessness occur

■ symptoms do not improve within 7 days or occur with fever

Questions or comments? 1-800-426-9391

Drug Facts

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

■ adults and children 12 years and over: take 1 tablet every 4 hours. Do not take more than 6 tablets in 24 hours.

■ children under 12 years: ask a doctor

Other information

■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

■ see end flap for expiration date and lot number

Inactive ingredients croscarmellose sodium, dextrose monohydrate, dibasic calcium phosphate dihydrate, F&D red #40, lecithin, magnesium stearate, maltodextrin, microcrystalline cellulose, silicon dioxide, sodium carboxymethylcellulose, sodium citrate dihydrate, titanium dioxide

Questions or comments? 1-800-426-9391

ORG0318-F
REV0518

Walgreens Pharmacist Survey
If this product is not manufactured or
distributed by Johnson & Johnson
Corporation, owner of the registered
trademark Sudafed PE® Day + Night Sinus
Congestion.

100% SATISFACTION
GUARANTEED

Walgreens
DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
ITEM 885140 W00000-0000-0
3 1191719998 6

PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

With any other product containing diphenhydramine, even one used on skin (Nighttime only)

Ask a doctor before use if you have heart disease, diabetes, high blood pressure, difficulty in urination due to enlargement of the prostate gland, breathing problem such as emphysema or chronic bronchitis (Nighttime only)

glaucoma (Nighttime only)

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers. (Nighttime only)

Drug Facts

Active ingredient (in each tablet)

Phenylephrine HCl 10 mg

Purpose

Nasal decongestant

Active ingredients (in each tablet)

Diphenhydramine HCl 25 mg

Phenylephrine HCl 10 mg

Purpose

Nasal decongestant

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- nasal congestion
- runny nose (Nighttime only)
- itchy, watery eyes (Nighttime only)
- itching of the nose or throat (Nighttime only)

temporarily relieves these symptoms due to the common cold:

- nasal congestion
- runny nose (Nighttime only)
- sneezing (Nighttime only)

temporarily relieves sinus congestion and pressure

Walgreens Pharmacist Recommended

Health expertise you rely on.™

B-2201-453485-09-H
REV0118A45348509

Walgreens

DAY & NIGHT PACK

NDC 0363-4385-09

DAYTIME • NON DROWSY

Wal-Phed® PE

SINUS CONGESTION

PHENYLEPHRINE HCl 10 mg / NASAL DECONGESTANT

MAXIMUM STRENGTH

- Relieves sinus pressure & congestion
- Pseudoephedrine free

12 TABLETS

NIGHTTIME

Wal-Phed® PE

SINUS CONGESTION

DIPHENHYDRAMINE HCl 25 mg / ANTIHISTAMINE
PHENYLEPHRINE HCl 10 mg / NASAL DECONGESTANT

- Relieves itchy, watery eyes, runny nose & itchy throat
- Pseudoephedrine free

8 TABLETS

Drug Facts (continued)**When using this product**

- do not exceed recommended dosage
- excitability may occur, especially in children (Nighttime only)
- marked drowsiness may occur (Nighttime only)
- alcohol, sedatives, and tranquilizers may increase drowsiness (Nighttime only)
- avoid alcoholic beverages (Nighttime only)
- use caution when driving a motor vehicle or operating machinery (Nighttime only)

Stop use and ask a doctor if

- symptoms do not improve within 7 days or occur with fever

Drug Facts (continued)**Directions**

- adults and children 12 years and over
 - take 1 tablet every 4 hours
 - do not take more than 6 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- each tablet contains: calcium 25 mg (Nighttime only)
- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C

Drug Facts (continued)**Inactive ingredients (Nighttime only)**

croscarmellose sodium, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, silicon dioxide, stearic acid, titanium dioxide, triacetin

Questions or comments? 1-800-426-9391

Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not give more than 6 doses in 24 hours unless directed by a doctor ■ mL = milliliter
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product

Age	Dose
children 6 to under 12 years	10 mL every 4 hours
children 4 to under 6 years	do not use unless directed by a doctor
children under 4 years of age	do not use

Other information ■ each 5 mL contains: sodium 3 mg

- store between 20-25°C (68-77°F). Do not refrigerate. ■ protect from light

Inactive ingredients acesulfame potassium, citric acid, EDTA disodium, FD&C blue #1, FD&C red #40, flavor, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING

DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED walgreens.com ©2018 Walgreen Co.

ITEM 357367 W10023-0218-F

PLD-H320D LB004646

ORG0617-F-R1

Walgreens

children's
Cold & Cough

DIPHENHYDRAMINE HCl /
ANTIHISTAMINE /
COUGH SUPPRESSANT
PHENYLEPHRINE HCl /
NASAL DECONGESTANT



Compare to Children's Triaminic®
Night Time Cold & Cough
active ingredients^{††}

- Nighttime
- Ages 6 to 11 years

4 FL OZ (118 mL)



**GRAPE
FLAVOR**

Drug Facts

Active ingredients (in each 5 mL)

Diphenhydramine HCl 6.25 mg.....Antihistamine/Cough suppressant
Phenylephrine HCl 2.5 mg.....Nasal decongestant

Purposes

Uses

- temporarily relieves
- sneezing ■ runny nose ■ itchy nose or throat
- itchy, watery eyes due to hay fever ■ nasal and sinus congestion
- cough due to minor throat and bronchial irritations as may occur with a cold

Warnings

Do not use ■ in a child under 4 years of age ■ in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product. ■ with any other product containing diphenhydramine, even one used on skin ■ for the purpose of making your child sleepy

Ask a doctor before use if the child has ■ heart disease ■ high blood pressure

- thyroid disease ■ diabetes ■ glaucoma ■ cough that occurs with too much phlegm (mucus)
- chronic cough that lasts, or as occurs with asthma ■ a breathing problem such as chronic bronchitis

Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers.

When using this product ■ do not exceed recommended dosage

- marked drowsiness may occur ■ excitability may occur, especially in children
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur

- symptoms do not improve within 7 days or occur with a fever ■ cough persists for more than 7 days, comes back, or occurs with a fever, rash, or persistent headache. These could be signs of a serious condition.

^{††}This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Children's Triaminic® Night Time Cold & Cough.

Drug Facts (continued)

- nervousness, dizziness or sleeplessness occurs
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- mL = milliliter
- shake well before using
- do not give more than 5 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- If needed, repeat dose every 4 hours while symptoms last
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

Weight (lb)	Age (yr)	Dose (mL)
48-95	6-11	10 mL
36-47	4-5	do not use unless directed by a doctor
under 36	under 4	do not use

Other information

- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients carboxymethylcellulose sodium, citric acid, D&C red #33, FD&C blue #1, FD&C red #40, flavor, glycerin, microcrystalline cellulose, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum

Questions or comments?
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

PARENTS:
Learn about drug medicine abuse
www.StopMedicineAbuse.org

DOSAGE CUP PROVIDED

"Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands."
"This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Plus Multi-Symptom Cold.
PLD-J253C FCO08009

NDC 0363-0034-04

Walgreens

Compare to the active ingredients in Children's Tylenol® Plus Multi-Symptom Cold*

MULTI-SYMPTOM Children's Cold + Flu

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
CHLORPHENIRAMINE MALEATE / ANTIHISTAMINE
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

- Relieves aches, fever, sore throat, sneezing, cough, runny & stuffy nose
- Oral suspension

AGES 6-11 YEARS

4 FL OZ (118 mL) Grape flavor

NDC 0363-0034-04

Walgreens

Compare to the active ingredients in Children's Tylenol® Plus Multi-Symptom Cold*

MULTI-SYMPTOM Children's Cold + Flu

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER
CHLORPHENIRAMINE MALEATE / ANTIHISTAMINE
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

- Relieves aches, fever, sore throat, sneezing, cough, runny & stuffy nose
- Oral suspension

AGES 6-11 YEARS

4 FL OZ (118 mL) Grape flavor

Drug Facts

Active Ingredients (in each 5 mL)

Active Ingredients (in each 5 mL)	Purposes
Acetaminophen 160 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 1 mg	Antihistamine
Dextromethorphan HBr 5 mg	Cough suppressant
Phenylephrine HCl 2.5 mg	Nasal decongestant

Uses

- temporarily relieves the following cold/flu symptoms
- minor aches and pains
- sore throat
- cough
- sneezing and runny nose
- headache
- nasal congestion
- stuffy nose
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses (10 mL) in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- to make a child sleepy
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has

- liver disease
- glaucoma
- thyroid disease
- diabetes
- heart disease
- high blood pressure
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

Ask a doctor or pharmacist before use if your child is taking

- the blood thinning drug warfarin
- sedatives or tranquilizers

When using this product

- do not exceed recommended dose (see Overdose warning)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- new symptoms occur
- redness or swelling is present
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

LOT No.:

Exp. Date:

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.

ITEM 510122 W10024-0322-L

3 11917 10357 0

Drug Facts (continued)

Age	Dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to 11 years	10 mL every 4 hours
children under 6 years	do not use

Other information

- each 10 mL contains: sodium 6 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive Ingredients citric acid, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sorbitol, sucralose

Questions or comments?
Call 1-877-763-3935 Monday-Friday 9AM-5PM EST

TAMPED EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING

DOSAGE CUP PROVIDED

*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
 **This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Children's Dimetapp® Cold & Allergy.

DISTRIBUTED BY: WALGREEN CO.
 200 WILMOT RD., DEERFIELD, IL 60015
Walgreens
100% SATISFACTION GUARANTEED
 walgreens.com
 ©2021 Walgreen Co.

NDC 0383-0312-08

Walgreens

Compare to the active ingredients in Children's Dimetapp® Cold & Allergy*

Children's Cold & Allergy Relief

BROMPHENIRAMINE MALEATE 2 mg / ANTIHISTAMINE PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

Alcohol Free

- Relieves itchy, watery eyes, runny nose, sneezing, stuffy nose & itching of the nose or throat

AGES 6 YEARS & OLDER

8 FL OZ (237 mL)

Grape flavor

NDC 0383-0312-08

Walgreens

Compare to the active ingredients in Children's Dimetapp® Cold & Allergy*

Children's Cold & Allergy Relief

BROMPHENIRAMINE MALEATE 2 mg / ANTIHISTAMINE PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

Alcohol Free

- Relieves itchy, watery eyes, runny nose, sneezing, stuffy nose & itching of the nose or throat

AGES 6 YEARS & OLDER

8 FL OZ (237 mL)

Grape flavor

Drug Facts

Active ingredients (in each 10 mL)

Brompheniramine maleate 2 mg.....Antihistamine

Phenylephrine HCl 5 mg.....Nasal decongestant

Purposes

Uses

- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis)
- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat
- temporarily restores free breathing through the nose

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

- taking any other oral nasal decongestant or stimulant
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter

ITEM 904372 W10024-0722-L



PLD-F2600 FC008257

Lot No.:
Exp. Date:

Drug Facts (continued)

Directions ■ do not take more than 6 doses in any 24 hour period
 ■ do not exceed recommended dosage ■ measure only with dosing cup provided. Do not use any other dosing device. ■ keep dosing cup with product ■ mL = milliliter

Age	Dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

Other information ■ each 10 mL contains: sodium 6 mg ■ store at 20-25°C (68-77°F) ■ do not refrigerate ■ protect from light

Inactive ingredients acesulfame potassium, anhydrous citric acid, EDTA disodium, FD&C blue #1, FD&C red #40, flavor, maltitol, propylene glycol, purified water, sodium benzoate, trisodium citrate dihydrate

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING

¹¹ This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Children's Dimetapp® Nighttime Cold & Congestion.

ITEM 904373 W10023-0717-F

PLD-C375A LB004650

ORG0617-F

3216
Walgreens

children's
COLD & CONGESTION
Wal-Tap®

DIPHENHYDRAMINE HCl 12.5 mg /
ANTIHISTAMINE / COUGH SUPPRESSANT
PHENYLEPHRINE HCl 5 mg /
NASAL DECONGESTANT

Compare-to Children's Dimetapp® Nighttime
Cold & Congestion active ingredients¹¹

- Nighttime
- Alcohol free
- 6 years & older

4 FL OZ (118 mL) GRAPE FLAVOR

NDC 0363-0058-04

Drug Facts**Active Ingredients (in each 10 mL)**

Diphenhydramine HCl 12.5 mg.....Antihistamine/Cough suppressant
 Phenylephrine HCl 5 mg.....Nasal decongestant


Uses ■ temporarily relieves these symptoms occurring with a cold, hay fever, or other upper respiratory allergies ■ nasal congestion ■ cough ■ runny nose
 ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings

Do not use ■ to sedate a child or to make a child sleepy ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ with any other product containing diphenhydramine, even one used on skin. **Ask a doctor before use if you have** ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ glaucoma ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema. **Ask a doctor or pharmacist before use if you are** ■ taking any other oral nasal decongestant or stimulant ■ taking sedatives or tranquilizers. **When using this product** ■ do not use more than directed ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children. **Stop use and ask a doctor if** ■ you get nervous, dizzy, or sleepless ■ symptoms do not get better within 7 days or occur with a fever ■ cough lasts more than 7 days, comes back, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015
 100% SATISFACTION GUARANTEED walgreens.com ©2017 Walgreen Co.



COMBO PACK		NDC 0363-9012-48
	<i>Compare to Vicks® DayQuil® & NyQuil® Cold & Flu LiquiCaps® active ingredients††</i>	
DAYTIME • NON-DROWSY	NIGHTTIME	
Cold & Flu	Cold & Flu	
ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT	ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER DEXTROMETHORPHAN HBr 15 mg / COUGH SUPPRESSANT DOXYLAMINE SUCCINATE 6.25 mg / ANTIHISTAMINE	
MULTI-SYMPTOM	MULTI-SYMPTOM	
		
ACTUAL SIZE	ACTUAL SIZE	
32 LIQUID CAPS	48 TOTAL LIQUID CAPS	16 LIQUID CAPS

Drug Facts (continued)

Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur
■ pain, nasal congestion, or cough gets worse, or lasts more than 5 days (children) or 7 days (adult) ■ fever gets worse, or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** **Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in any 24-hour period ■ measure only with dosing cup provided. Do not use any other dosing device. ■ mL = milliliter ■ keep dosing cup with product

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information ■ each 15 mL contains: sodium 12 mg
■ store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

¹Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
^{††}This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

DISTRIBUTED BY: WALGREEN CO. 200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED walgreens.com ©2022 Walgreen Co.

ITEM 271558 W10369-0922-F



W10369-0922-F
REV-0922

PARENTS:
Learn about your medicine online
www.StopTheMedicineHouse.org
PLD B3800 LB000867

Walgreens

Compare to the active ingredients
in Vicks® DayQuil® Cold & Flu^{††}



DAYTIME • NON-DROWSY Cold & Flu

ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

Multi-Symptom

- Relieves aches, fever & sore throat, cough & nasal congestion
- For ages 6 years & over
- Alcohol free
- Antihistamine free

12 FL OZ (355 mL)

NDC 0363-4661-12

Drug Facts

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

Active ingredients (in each 15 mL)

Active ingredients (in each 15 mL)	Purposes
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses ■ temporarily relieves common cold and flu symptoms
■ minor aches and pains ■ headache ■ fever ■ sore throat
■ nasal congestion ■ cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if: ■ adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount
■ child takes more than 4 doses (15 mL each) in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ liver disease
■ heart disease ■ high blood pressure ■ thyroid disease
■ diabetes ■ a sodium-restricted diet
■ trouble urinating due to an enlarged prostate gland
■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dosage.

3220

NDC 0363-9974-08

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL
AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING**Drug Facts****Active ingredients (in each 15 mL) Purposes**

Dextromethorphan HBr 10 mg..... Cough suppressant
Phenylephrine HCl 5 mg..... Nasal decongestant

Uses

■ temporarily relieves common cold and flu symptoms
■ minor aches and pains ■ headache ■ sore throat
■ nasal congestion ■ fever
■ cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if: ■ adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount ■ child takes more than 4 doses (15 mL each) in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

← PEEL HERE **Drug Facts** (continued under label)

Walgreens

Compare to the active ingredients
in Vicks® DayQuil® Cold & Flu*



DAYTIME • NON-DROWSY Cold & Flu

**ACETAMINOPHEN 325 mg / PAIN RELIEVER /
FEVER REDUCER**

**DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT**

Multi-Symptom

- Relieves aches, fever, sore throat, cough & nasal congestion
- For ages 6 years & over
- Antihistamine free
- Alcohol free

8 FL OZ (237 mL)

PARENTS:

Learn about teen medicine abuse
www.StopMedicineAbuse.org

*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

DISTRIBUTED BY:

WALGREEN CO.
200 WILMOT RD.,
DEERFIELD, IL 60015
100% SATISFACTION
GUARANTEED
walgreens.com

©2022 Walgreen Co.

PLD-B380D LB009015

W3ORG1022-F
REV-0323

how2recycle.info
PLASTIC BOTTLE
PLASTIC CUP

ITEM 816520 W11042-0423-F



3 11917 12396 7

Drug Facts (continued)

■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ liver disease
■ heart disease ■ high blood pressure ■ thyroid disease
■ diabetes ■ a sodium-restricted diet ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dosage.

Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur ■ pain, nasal congestion, or cough gets worse, or lasts more than 5 days (children) or 7 days (adult) ■ fever gets worse, or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended

Drug Facts (continued)

dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in any 24-hour period ■ measure only with dosing cup provided. Do not use any other dosing device.
■ keep dosing cup with product ■ mL = milliliter

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

■ each 15 mL contains: sodium 12 mg
■ store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients: citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call 1-877-753-3636 Monday-Friday 9AM-6PM EST

Walgreens

Cold, Flu & Sore Throat

Walgreens

Compare to Maximum Strength Mucinex® Fast-Max® Cold, Flu & Sore Throat active ingredients*

NDC 0083-1171-16

Cold, Flu & Sore Throat

ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER
 DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT
 GUAIFENESIN 200 mg / EXPECTORANT
 PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

MAXIMUM STRENGTH

ALCOHOL FREE

- Relieves fever, sore throat, chest congestion & nasal congestion
- Controls cough
- Thins & loosens mucus
- 12 years & older

16 SOFTGELS*
 (*LIQUID-FILLED CAPSULES)



ACTUAL SIZE

PHARMACY

Walgreens
 Trusted since 1901™

RECOMMENDED

Health expertise
 you rely on™

Drug Facts

Active ingredients

(in each softgel)

Acetaminophen 325 mg... Pain reliever/fever reducer
 Dextromethorphan HBr 10 mg... Cough suppressant
 Guaifenesin 200 mg... Expectorant
 Phenylephrine HCl 5 mg... Nasal decongestant

Purposes

Uses

- temporarily relieves these common cold and flu symptoms:
 - headache
 - nasal congestion
 - sore throat
 - minor aches and pains
 - cough
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:
 - more than 12 softgels in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks daily while using this product

- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
 - skin reddening
 - blistering
 - itching
 - skin peeling

- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease

Drug Facts (continued)

- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- new rashes, diarrhea or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see OVERDOSE WARNING)
- do not take more than 12 softgels in any 24-hour period
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use

Other information

- store between 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

FD&C Yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide

Questions or comments?

Call 1-888-353-8782

Walgreens Pharmacist Survey
 This product is not manufactured or distributed by Walgreen Co. owner of the registered trademarks Maximum Strength Mucinex® Fast-Max® Cold, Flu & Sore Throat.

DISTRIBUTED BY:
 WALGREEN CO.
 200 WILMOT ROAD
 DEERFIELD, IL 60015

Walgreens
100% SATISFACTION GUARANTEED
 walgreens.com
 ©2019 Walgreen Co.
 MADE IN INDIA

ORQ0619-F

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND TAMPER-EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.

ITEM 513230 W00000-0000-0



LOT NO:
 EXP DATE:

Walgreens

Compare to Maximum Strength
Mucinex® Fast-Max® Cold, Flu &
Sore Throat active ingredients††

NDC 0363-9016-16

Cold, Flu & Sore Throat

ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT
GUAIFENESIN 200 mg / EXPECTORANT
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

MAXIMUM STRENGTH

ALCOHOL FREE

- Relieves fever, sore throat, chest congestion
& nasal congestion
- Controls cough
- Thins & loosens mucus
- 12 years & older

16 SOFTGELS*
(*LIQUID-FILLED CAPSULES)



ACTUAL SIZE